





## European XFEL Hibrid Undulatörleri'nin İşletime Hazır Hale Getirilmesi ve XFEL'lerin Yüksek Teknoloji Uygulamaları

Doç. Dr. Bora KETENOĞLU

Ankara Üniversitesi Mühendislik Fakültesi

Fizik Mühendisliği Bölümü


[bketen@eng.ankara.edu.tr](mailto:bketen@eng.ankara.edu.tr)



# European XFEL Bilimsel Araştırma Merkezi Hamburg, Almanya



<https://www.xfel.eu>






**İki uçtaki tünel derinlikleri:** 6 - 38 metre  
**Hızlandırıcı:** 1.7 km uzunluğunda elektron linak  
Elektron demet enerjisi 17.5 GeV  
(20 GeV'e kadar artırılabilir)  
**ID'ler:** Toplam 91 adet U40 ve U68 undulatörler  
"5m uzunluğunda ve 8 ton ağırlığında"






# European XFEL'in Dünya'daki diğer X-Işını Serbest Elektron Lazeri Tesisleri ile Karşılaştırması





# Almanya/Hamburg'da Çalışmakta Olan 3.4 km Uzunluklu X-Işını Dalgaboylu Serbest Elektron Lazeri Tesisi






| Planlanan Bilimsel Deneyler                                                                                                                                                                       | nm Ölçekli Yapılar                                                                    | Ultrahızlı Süreçler                                                                   | Sıradışı Durumlar                                                                     | Demet Hattı |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|-------------|
| SPB (Tek Parçacıklar, Yığınlar & Biyomoleküller)<br>Ultrahızlı koherent kırınım görüntüleme, Tek parçacıkların yapı belirlenmesi: atomik yığınlar, biyomoleküller, virüs parçacıkları ve hücreler |    |    |                                                                                       | SASE1       |
| FXE (Femtosaniye X-ışını Deneyleri) Katıların, sıvıların ve gazların dinamiklerinin zaman çözünürlüklü incelemeleri                                                                               |                                                                                       |    |                                                                                       | SASE1       |
| MID (Malzeme Görüntüleme ve Dinamikleri)<br>Nano aygıtların yapı belirlenmesi ve nano boyutta dinamikler                                                                                          |    |    |                                                                                       | SASE2       |
| HED (Yüksek Enerji Yoğunluklu Madde Deneyleri)<br>Sert X-ışını dalga boylu SEL ile maddenin sıradışı durumlarda incelenmesi, örneğin: yoğun plazmanın algılanması                                 |                                                                                       |    |    | SASE2       |
| SQS (Küçük Kuantum Sistemleri) Şiddetli alanlarda atomların, iyonların, moleküllerin ve yığınların incelenmesi, Doğrusal olmayan fenomen                                                          |  |  |  | SASE3       |
| SCS (Spektroskopi ve Koherent Saçılma) Elektronik ve atomik yapı ve yumuşak X-ışınları ile nanosistemlerin dinamiği ve tekrarlanamayan biyolojik nesneler                                         |  |  |                                                                                       | SASE3       |

# Doğrusal Elektron Hızlandırıcısı









**$\sim 10^{10}$  elektron içeren elipsoid şeklindeki paketçikler, ışık hızına çok yakın hızlara kadar ( $v \approx c$ ) hızlandırılırlar.**

# Planar Undulator (Düzlemsel Salındırıcı)






# 5m Uzunluęunda U40 Undulat6r6 (Birbirinden Baęımsız 4 Ayrı Motor ile Kontrol)








# European XFEL Undulator Sistemleri Manyetik Ölçüm Laboratuvarı (21 °C Sabit Sıcaklıkta)

6.5 m Uzunluğunda  
Manyetik Ölçüm Düzenek














**Undulatörlerin bütün kutupları manyetik labda başarılı bir şekilde ayarlanıp landmark ölçümleri yapıldıktan sonra, tünele transfer edilmeye hazır hale gelirler.**







# **X-ışını Dalgaboylu Serbest Elektron Lazerleri'nin Disiplinlerarası Kullanım Alanları**



# **Fizik**'te **X-ışını Serbest Elektron Lazeri Deneyleri**

**Katıların, sıvıların, gazların statik özellikleri**



**Elektrik veya magnetik alan atmaları yada ultrakısa optik lazer flaşları ile başlatılan dinamik süreçler**

**Optik ışık atmaları ile tetiklenen faz geçişleri ve titreşimleri süreçleri**

**Dayanıklılık ve elastiklik gibi geliştirilmiş özellikli yeni malzemeler için kristallenme veya çatlak oluşumu ve yayılımı süreçleri**




# Biyoloji'de X-ışını Serbest Elektron Lazeri Deneyleri



**XSEL atması bir biyomolekül ile etkileştiğinde, saçılan X-ışınları kırınım deseni oluşturur. Farklı farklı hizalanmış molekülden saçılan pekçok desen yardımı ile, biyomolekülün üç boyutlu yapısı belirlenir.**





# Kimya'da X-ışını Serbest Elektron Lazeri Deneyleri



## Moleküler filmler:

Bir kimyasal reaksiyon, geleneksel lazer flaşı ile tetiklenir. Ardından, molekülde oluşan değişikliklerin fotoğraf karelerini çekmek üzere, değişken zaman aralıklarıyla SEL atmaları gönderilir.

# **Astrofizik'te ve Sıradışı Durumlar Bilimi'nde X-ışını Serbest Elektron Lazeri Deneyleri**




**Yıldızların ve  
gezegenlerin  
plazma ve  
diğer sıradışı  
durumlarının  
incelenmesi**




# Malzeme Bilimi'nde X-ışını Serbest Elektron Lazeri Deneyleri



Uçak yapımında  
kullanılan  
kaynaklı dikişlerin  
kalite kontrol  
muayenesinde.



# Elektronik'te X-ışını Serbest Elektron Lazeri Deneyleri




**Bilgisayarların ve diğer ticari elektronik cihazların veri aktarım hızı ve depolama kapasitelerinin geliştirilmesi.**



# Çevre ve Enerji'de

## X-ışını Serbest Elektron Lazeri Deneyleri

Güneş pillerinde meydana gelen süreçlerin atomik düzeyde çalışılması.



# **Yaşam Bilimleri Uygulamaları**



# Biyofizik ve Moleküler Biyoloji:

## XFEL'leri Kullanarak Biyo – “Kristalografi ve Spektroskopi”



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

**ScienceDirect**

Current Opinion in  
**Structural Biology**

**Metalloprotein structures at ambient conditions and in real-time: biological crystallography and spectroscopy using X-ray free electron lasers**


Jan Kern<sup>1,2</sup>, Vittal K Yachandra<sup>1</sup> and Junko Yano<sup>1</sup>



**Current Opinion in Structural Biology (2015) 34, 87-98.**

e) Fotosistem II'nin  
katalitik  
merkezlerindeki  
atomların  
pozisyonları.

Mavi küreler SR  
deneylerinden elde  
edilen pozisyonları,  
diğer renkler ise cryo-  
Serial  
Femtosecond  
Crystallography  
(SFX)  
pozisyonlarını temsil  
etmektedir.




Goniometer based crystallography experiments of metalloenzymes at XFEL sources. (a) Schematic of the experimental approach using a large crystal to collect still shots in a defined orientation with respect to each other, translating by 50  $\mu\text{m}$  between shots and rotating by a small angle [11\*\*]. (b) Alternatively many small crystals mounted in a mesh or similarly can be used. The few  $\mu\text{m}$  focused beam leaves imprints on the sample [10\*]. (c) Omit electron density for the heme group in myoglobin obtained from measuring diffraction of several hundred individual small crystals [10\*]. (d) 2Fo - Fc (blue, contoured at 1.0 $\sigma$ ) and Fo - Fc (red, contoured at 3.0 $\sigma$ ) difference density maps calculated for [FeFe] hydrogenase at the site of one of the Fe-S clusters obtained from measurements on five large crystals [10\*]. (e) Positions of the atoms in the catalytic center of PS II as obtained from cryo SFX is shown in comparison to the positions obtained by SR experiments (blue spheres) [9\*\*]. (f) Radiation damage free omit map for the FeCu center of CoC showing the position of the peroxide ligand [11\*\*].




## b) Liquid Jet düzenegi ile forward scattering şematik olarak gösterilmiştir.

Bu düzenek, tetiklemeli  
optik lazer atmaları ve  
spektroskopik  
ölçümlerle  
birleştirilebilir (örneğin,  
X-ray Emission  
Spectroscopy gibi).



Jet based crystallography experiments of metalloenzymes at XFEL sources. (a) Examples of nano- to micrometer size crystals used, e.g. thermolysin [17], PS II [36\*\*] and toxin crystals (grown inside cells) [29]. (b) Schematic of the experimental approach using a liquid jet for sample delivery to the interaction point and collecting the forward scattering is shown. This setup can be combined with triggering by optical laser pulses and with spectroscopic measurements, e.g. X-ray emission spectroscopy. XFEL structures obtained: (c) PS I with electron density shown in blue and omit maps for the  $Fe_4S_4$  clusters in red [25]. (d) Electron density for one of the  $Fe_4S_4$  clusters in ferredoxin, with omit map shown in green [32]. (e) PS II with overall structure of one monomer (bottom left), the oxygen evolving complex (bottom right) [36\*\*] and the anomalous difference density from the Mn atoms in the OEC (top right) [31\*]. (f) PBRC with overall structure and omit maps for the cofactors (left) and detail for the electron density obtained for one of the heme groups (right) [33].




## a) Proteinlerdeki ve enzimlerdeki dinamik süreçler için zaman mertebeleri

Titreşimsel hareketler ve exciton transferleri için femtosecond deneyler gerekli.

Time resolved experiments using fs X-ray pulses at the CXI instrument at LCLS. (a) Timescales for dynamic processes in proteins/enzymes. (b) Time resolved WAXS measurements on the photosynthetic bacterial reaction center. Upon light excitation a ultrafast protein movements in the ps time regime was visible [44\*]. (c) Illustration of the protein-quake observed within a few ps in myoglobin upon photodissociation of CO [45\*], red shaded areas are involved in the movement leading to a volume increase of  $\sim 200 \text{ \AA}^3$ . Time resolved serial femtosecond crystallography on photoactive yellow protein, showing the electron density for (d) the dark state as well as difference electron density for (e) data recorded 10 ns after light excitation and (f) 1 μs after light excitation. A movement of the S-atom of the chromophore is visible already in the 10 ns data (Adapted from [46]). (g) X-ray emission spectra of Mn in photosystem II. Spectra are shown for the 2F state (S<sub>2</sub>), as well as for an intermediate 250 μs after the 3<sup>rd</sup> light flash and 0.5 s after the 3<sup>rd</sup> light flash (S<sub>0</sub>-state). The difference spectra (middle panel) indicate a reduction in the S<sub>2</sub>-S<sub>0</sub> transition (as expected from SR measurements — bottom panel), but no specific reduction is visible for the 250 μs state [31\*].



## c) Mn<sup>2+</sup> ve MnCl<sub>2</sub> 1s-3p orbitalleri arasındaki “K<sub>β</sub> emission” enerjileri için Von Hamos Spectrometer



X-ray spectroscopic measurements using XFEL pulses. (a) Energy level diagram illustrating the transitions involved in L- and K-edge transition metal spectroscopy. (b) Examples for Fe K-edge absorption spectra showing transient changes in spin state [53]. (c) Experimental setup for collecting K $\beta$  emission spectra using a dispersive Von Hamos spectrometer and a sample injector. In addition spectra of two Mn compounds collected at LCLS are shown [51]. (d) Experimental setup for collection of Mn L-edge spectra at an XFEL from dilute aqueous solutions of Mn compounds using a reflective zone plate (RZP) spectrometer. The spectrum shown was recorded at LCLS using 500 mM MnCl<sub>2</sub> in water [56].


# **Yüksek Teknoloji Uygulamaları**



# Doğadaki Çok Hızlı Süreçlerin İncelenmesi



- Charge transfer in catalytic/photovoltaic systems – 1 fs and longer
- Phase change in materials – 2 fs and longer
- Ultrafast spintronics – fs and longer
- Control electron-ion motions in chemical reactions – 1 fs and longer
- X-ray induced processes – 50 as and longer
- Strong field physics – zeptoseconds and longer

**Electron  
dynamics**






**Image charge transfer in complex systems relevant to energy, catalysis using coherent x-ray spectroscopy spanning many elemental absorption edges simultaneously**



# Yüksek Harmoniklerle Nanodinamiklerin Görüntülenmesi




**Capture charge-spin-phonon dynamics at multiple sites:** (*Nature* **471**, 490 (2011), *PNAS* **109**, 4792 (2012); *Nature Comm* **3**, 1037 (2012); *Nature Comm* **3**, 1069 (2012))



**Coupled electron-nuclear dynamics in molecules:** (*Science* **317**, 1374 (2007), *Science* **322**, 1081 (2008), *Nature Phys.* **8**, 232 (2012), *PRL* **109**, 073004 (2012))




**Nanoscale imaging:** Record tabletop 22nm resolution (*Op. Ex.* **19**, 22470 ('11); **17**, 19050 ('12); *Nature* **463**, 214 (2010))



**Nanoscale energy transport:** probe nanoscale energy/strain flow (*Nature Materials* **9**, 26 (2010); *Nano Letters* **11**, 4126 (2011); *PRB* **85**, 195431 (2012))



## 20. Yüzyılın İkinci Yarısından Günümüze Veri Depolama Evrimi





## The Nobel Prize in Physics 2007

"for the discovery of Giant Magnetoresistance"



Photo: U. Montan

**Albert Fert**




Photo: U. Montan

**Peter Grünberg**

*This year's Nobel Prize in Physics is awarded to **ALBERT FERT** and **PETER GRÜNBERG** for their discovery of Giant Magnetoresistance. Applications of this phenomenon have revolutionized techniques for retrieving data from hard disks. The discovery also plays a major role in various magnetic sensors as well as for the development of a new generation of electronics. The use of Giant Magnetoresistance can be regarded as one of the first major applications of nanotechnology.*

### Better read-out heads for pocket-size devices


Constantly diminishing electronics have become a matter of course in today's IT-world. The yearly addition to the market of ever more powerful and lighter computers is something we have all started to take for granted. In particular, hard disks have shrunk – the bulky box under your desk will soon be history when the same amount of data can just as easily be stored in a slender laptop. And with a music player in the pocket of each and everyone, few still stop to think about how many cds' worth of music its tiny hard disk can actually hold. Recently, the maximum storage capacity of hard disks for home use has soared to a terabyte (a thousand billion bytes).



A nanometre is one millionth of a millimetre and corresponds to only a few layers of atoms. Albert Fert's and Peter Grünberg's original structures for giant magnetoresistance consisted of only a few layers of atoms of magnetic iron (red) sandwiched with chromium (grey).



# SONUÇ



## 21. Yüzyıl'ın Bilim ve Teknolojisine Yön Veren “X-Işını Dalgaboylu FEL” Tesisleri

*İlginiz ve Katılımınız için  
Teşekkür Ederim.*